

Neck Pain, Cervicalgia

Diagnosis/Condition:	Neck Pain, Cervicalgia Cervical sprain
Discipline:	Integrated
ICD-10 Codes:	M54.2; M53.1; M99.01; S13.4XXA S13.8XXA
Origination Date:	1996
Review/Revised Date:	07/2023
Next Review Date:	07/2025

Neck pain is second only to low back pain as the most common musculoskeletal disorder in population surveys and in primary care. Depending on the case definition of “neck pain,” from 30 to 50% of the adult population experiences neck pain each year.¹ Neck pain is common in children as well, with prevalence estimates of 40% of pre-adolescents and teenagers.^{2,3}

Classification of cervicalgia according to ICD-10 includes neck pain (M53.2), cervicobrachial syndrome (M53.1), and biomechanical lesions (M99.01). Strictly speaking, cervicalgia is neck pain without radiation, although radiating symptoms are commonly encountered.

Other neck pain classification schemes consider acute, subacute, and chronic pain problems based on the time course of the disorder. Acute refers to the first month, subacute 2-3 months and chronic >3 months. Subacute also incorporates intermittent recurrences of pain interrupted by pain free intervals. Yet another classification scheme has been proposed by the Bone and Joint Decade Taskforce on Neck Pain (see assessment section below).⁴ The Task Force on Neck Pain aims to assess the literature to best determine the epidemiology, diagnosis, prognosis, economic costs, and treatment of neck pain.⁵ Task Force recommends manual approaches (including chiropractic and acupuncture) as first-line therapies for the treatment of chronic mechanical neck pain without radiculopathy.

Estimates of the global burden of disease attributed to neck pain indicate neck pain is the 4th highest condition causing disability.⁶ Neck and back pain combined cost an estimated \$86 billion in the US. Yet despite its prevalence, high disability, and health care costs, research devoted specifically to neck pain is minimal compared to other spinal problems such as cervical radiculopathy, low back pain, and lumbar radiculopathy.

Neck pain is frequently reported to and treated by integrative healthcare (IH) providers. US surveys indicate that neck pain is one of the most common complaints for which individuals seek treatment with Complementary and Integrative Health (CIH) therapies.^{7,8} There is some evidence that manual therapy and acupuncture techniques have been found to be less costly

and more effective than either physical therapy or general practitioner care in improving neck pain.^{9,10}

While most individuals with acute neck pain do not seek health care, those that do account for a disproportionate amount of health care costs. Most neck pain is not attributable to a specific disease or disorder and is labeled as muscular, mechanical, or postural neck pain. Despite decades of research and posturing to explain chronic neck pain based on a specific disease or injury, and despite increasingly sophisticated diagnostic imaging assessment, little advance has been made in achieving a specific structural diagnosis for this prevalent condition. Individuals with neck pain do not typically experience full remission; it is estimated that 50-80% will re-experience symptoms in 1-5 years.^{11,12}

Subjective Findings and History

- Risk factors for neck pain include age, gender, and heredity.¹
- Modifiable risk factors include tobacco exposure (smoking and second hand), low social support at work, high quantitative work demands, sedentary, precision, and repetitive work.¹
- Traumatic neck pain is the onset of pain and paraspinal muscle spasm begins either immediately after an injury (e.g., motor vehicle collision, sports injury) or gradually over the succeeding 24 hours.
- Nontraumatic neck pain comes on as a result of minor traumatic events not singularly capable of producing injury such as postural strain associated with cell phone use (“text neck”), acute myogenic torticollis (wry neck), and other overuse disorders.
- Local pain, sometimes accompanied by referred pain, diffuse (scleratogenous pain distribution).
- Loss of flexibility.
- Pain is usually relieved by rest and aggravated by motion.

Objective Findings

The validity of most commonly used objective tests is lacking. There is support for subjective self-report assessment in monitoring the patient’s course and response to treatment.

- Postural evaluation may reveal asymmetry, misalignment or decrease of normal spinal curvature, and may present with lateral list.
- Decrease/loss of normal cervical ROM may be present.
- Palpation may reveal: Segmental joint dysfunction/subluxation, tenderness with pressure over involved tissues, muscle spasm or tautness of paravertebral muscles, myofascial trigger points, and tenderness of acupuncture points.
- Orthopedic and neurological examination directed at differentiating neurogenic from other sources of pain: absence of nerve compression signs (e.g., absence of muscle weakness); orthopedic tests may reproduce the pain (e.g., foraminal compression and other tests that cause spinal motion may increase neck pain).

- Radiographic examination: depending on age and history of prior episodes (see CHP's Radiographic Guidelines behind the provider log in at www.chpgroup.com: *Providers* tab, *Provider Hub* tab, *Provider Log-in*, *Clinical Tools* tab).
- Pulse and tongue diagnosis findings will vary.

Assessment

- Rule out “red flags” of serious pathology especially in patients seeking care for neck pain without frank trauma. Serious disorders to rule out include pathologic fractures, neoplasm, systemic inflammatory diseases, infections, cervical myelopathy and/or previous cervical spine or neck surgery, or open injury.¹³
- The clinical impression should indicate the specific anatomical structures and acupuncture channels involved, and clinically correlate them with the mechanism of injury, history, subjective complaints, and objective findings.
- Risk factors for chronicity: Significant trauma, co-morbidity (degenerative disc disease, segmental instability, osteoporosis, spine deformity), age, socio-economic factors.

The Task Force on Neck Pain recommends a 4-grade classification system of neck pain severity.¹¹

- Grade I neck pain: No signs or symptoms suggestive of major structural pathology and no or minor interference with activities of daily living; will likely respond to minimal intervention such as reassurance and pain control; does not require intensive investigations or ongoing treatment.
- Grade II neck pain: No signs or symptoms of major structural pathology, but major interference with activities of daily living; requires pain relief and early activation/intervention aimed at preventing long-term disability.
- Grade III neck pain: No signs or symptoms of major structural pathology, but presence of neurologic signs such as decreased deep tendon reflexes, weakness, and/or sensory deficits; might require investigation and, occasionally more invasive treatments.
- Grade IV neck pain: Signs or symptoms of major structural pathology, such as fracture, myelopathy, neoplasm, or systemic disease; requires prompt investigation and treatment.

TCM Perspective

According to TCM theory, neck pain corresponds to a type of obstruction in the circulation of qi and blood in the channels and collaterals, traditionally known as ‘Bi syndrome.’ This syndrome indicates pain, soreness or numbness of the muscles, tendons and joints from invasion of ‘external pathogenic factors’, i.e., wind, cold and/or dampness, or internal imbalance. These pathogenic factors obstruct the circulation of qi and blood in the channels and collaterals, causing pain.¹⁴

Plan

Following a trauma (e.g., whiplash), patients should be encouraged to resume pre-accident activity as quickly and safely as possible. Patients should also be reassured that neck pain

normally has a good prognosis and regaining normal function is highly likely. Interventions that focus on regaining function and returning to work as soon as possible are relatively more effective than interventions that do not have such a focus.

Chiropractic Adjustments/Naturopathic Manipulation:

- Cervical and thoracic manipulation, cervical mobilization.^{15,16,,17,18,19,20}
- Cervical posture corrective orthotic device.²¹
- There is low level evidence that intermittent cervical traction is helpful for neck pain.²²
- A 2014 study provides strong recommendations for treatment of chronic neck pain with manipulation, manual therapy and exercise in combination with other modalities²³
- Treatment of seniors (65 years or older) with neck pain using spinal manipulative therapy proved beneficial when combined with supervised and unsupervised home exercise with the supervised component adding little benefit.²⁴

Soft Tissue Therapies:

- Physical therapy modalities, including manual therapy with a multi-modal approach.^{25,26, 27}
- Manual pressure.²⁸
- Passive and active soft tissue therapy.²⁹
- Massage Therapy.³⁰
- Low level laser therapy may be useful in chronic neck pain.³¹

Movement and Exercise:

- Alexander Technique.³²

Over-The-Counter Drugs:

- Analgesics, NSAIDS, muscle relaxants. There is insufficient data to suggest that one medication is superior or more effective to any other or to non-pharmaceutical interventions.³³ They also demonstrate no observable differences in short or long-term assessment of Quality of Life Years (QALYs).^{11,33}
- Topical salicylic acid, capsaicin, CBD, and other topical analgesic creams.

Nutritional Supplements:

- Oral and intramuscular injections of magnesium; bromelain and other proteolytic enzymes; Vit. C and B-complex.

Homeopathy:

- Bryonia, Arnica, Rhus tox, Mag phos, Calc phos et al as indicated by symptomatology.

Western Herbal Medicine:

- Botanical medicines: Curcumin, Devil's club, Valerian topically, Cimicifuga, nervines for sleep and relaxation, Arnica, DMSO.
- Topical treatments (e.g., comfrey poultice, hypericum, arnica, wintergreen).

- Botanical supplementation to reduce inflammation (*Curcuma longa* (turmeric), *Capsicum annuum* (cayenne), *Arnica montana* (arnica), *Ruta graveolens* (rue), *Hypericum perforatum* (St. John's wort), and *Gaultheria procumbens* (wintergreen)).

Immobilization, Bracing, Taping:

- Braces/supports.
- Posture training, ergonomic evaluation, educational interventions, for example cranio-cervical flexion exercises.³⁴
- Activity/work restrictions and rest, if appropriate. These may include: No repetitive motion, lifting, grasping, pinching.

Acupuncture:

While no RCTs have been conducted on the benefits of acupuncture for *acute* neck pain, approximately 20 RCTs have assessed the value of acupuncture for the treatment of *chronic* neck pain. With the exception of five trials (n≥100)^{35,36,37,38,39}, the published RCTs have been summarized in four systematic reviews^{40,41,42,43}. Based on the literature the following conclusions can be drawn:

The evidence supports the benefit of acupuncture for pain relief in chronic neck pain.

- The most recent systematic reviews (2006 & 2009) demonstrate the effectiveness of acupuncture adjunctive to usual medical care for pain, function, and disability at the end of treatment and at short-term follow-up.^{44,45}
 - This is supported by a new RCT (2011) not included in the above reviews.⁴⁶
- An RCT (2015) suggests acupuncture provides long-term benefit (1-year follow-up) when used in addition to usual care.⁴⁷
 - This is supported by the findings from individual patient data meta-analyses (n=3; 357 participants); suggests acupuncture is an effective treatment for chronic neck pain and is superior to sham controls.⁴⁸

The evidence supports the cost-effectiveness of acupuncture for the treatment of chronic neck pain.

- A large-scale trial (n=3,451) demonstrated cost-effectiveness at 1 and 4 -years.⁴⁹

Lifestyle Counseling:

- Psychoemotional counseling.⁵⁰
- Cognitive and Mind/Body Therapies.⁵¹

Mind-Body Therapies:

- Yoga.^{52,53}
- QiGong:
 - A systematic review (N=5; 399 participants) suggests benefit.⁵⁴

Outcome Assessment Tools

- Visual analog pain scale (VAS), numeric pain rating scale (NRS), quadruple VAS and NRS.

- Neck pain disability index (NDI).
- Patient Specific Functional Scale.
- MYMOP2 (Make Yourself Medical Outcome Profile version 2).⁵⁵
- QALY (Quality of Life Years).⁵⁶
- Northwick Park Questionnaire.⁵⁷

Referral Criteria

Referral to an appropriate specialist may be appropriate after 4-6 weeks of care without symptomatic or functional improvement or upon onset of (progressive) neurologic deficit.

Resources for Clinicians

Haldeman S, et.al. A Best Evidence Synthesis on Neck Pain: Findings from The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*. February 15, 2008, Volume 33, Issue 4S Supplement.

Martel JW, Potter SB. Evaluation and Management of Neck and Back Pain. *Semin Neurol*. 2019;39(1):41-52.

McDonald J, Janz S. The Acupuncture Evidence Project: A Comparative Literature Review (Revised Edition). Brisbane: Australian Acupuncture and Chinese Medicine Association Ltd; 2017. <http://www.acupuncture.org.au/> (accessed June 2021)

Society for Acupuncture Research. Acupuncture for the Treatment of Cervical Neck Pain. July 2021. www.AcupunctureResearch.org. (accessed June 2021)

Resources for Patients

Spine-health.com publishes original, award-winning articles written for patients by over 80 physician authors and peer-reviewed by a 23-member Medical Advisory Board. This trusted, independent site is supported by hundreds of physician members and visited by millions of patients and their physicians. <http://www.spine-health.com/pain/neck-pain-0?page=1>

MedlinePlus will direct you to information to help answer health questions. MedlinePlus brings together authoritative information from NLM, the National Institutes of Health (NIH), and other government agencies and health-related organizations. <http://www.nlm.nih.gov/medlineplus/ency/article/003025.htm>

Clinical Pathway Feedback

CHP desires to keep our clinical pathways customarily updated. If you wish to provide additional input, please use the e-mail address listed below and identify which clinical pathway you are referencing. Thank you for taking the time to give us your comments.

Clinical Services Department: cs@chpgroup.com

Disclaimer Notice

The CHP Group (CHP) Clinical Pathways are a resource to assist clinicians and are not intended to be nor should they be construed/used as medical advice. The pathways contain information that may be helpful for clinicians and their patients to make informed clinical decisions, but they cannot account for all clinical circumstances. Each patient presents with specific clinical circumstances and values requiring individualized care which may warrant adaptation from the pathway. Treatment decisions are made collaboratively by patients and their practitioner after an assessment of the clinical condition, consideration of options for treatment, any material risk, and an opportunity for the patient to ask any questions.

CHP makes no representation and accepts no liability with respect to the content of any external information cited or relied upon in the pathways. The presence of a particular procedure or treatment modality in a clinical pathway does not constitute a representation or warranty that this service is covered by a patient's benefit plan. The patient's benefit plan determines coverage.

¹ Hogg-Johnson S, et al. The burden and determinants of neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *J Manipulative Physiol Ther.* 2009 Feb;32(2 Suppl):S46-60.

² Weber Hellstenius S. Recurrent neck pain and headaches in preadolescents associated with mechanical, dysfunction of the cervical spine: A cross-sectional observational study with 131 students. *Journal Of Manipulative And Physiological Therapeutics.* October 1, 2009;32(8):625.

³ Ståhl M, et al. Neck pain in adolescence. A 4-year follow-up of pain-free preadolescents. *Pain.* July 2004; 110 (1): 427-431.

⁴ Haldeman S, Carroll L, Cassidy JD, Schubert J, Nygren A, Cn B, Joint Decade – Task Force on Neck P, Its Associated D: The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: executive summary. *Spine (Phila Pa 1976)* 2008, 33(4 Suppl):S5-7.

⁵ Hurwitz, E.L., et al., Treatment of neck pain: noninvasive interventions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)*, 2008. 33(4 Suppl): p. S123-52.

⁶ Hoy D, March L, Woolf A, et al The global burden of neck pain: estimates from the Global Burden of Disease 2010 study *Annals of the Rheumatic Diseases* 2014;73:1309-1315.

⁷ Eisenberg, D.M., et al., Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *JAMA*, 1998. 280(18): p. 1569-1575.

⁸ Barnes, P.M., B. Bloom, and R.L. Nahin, Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report*, 2008(12): p. 1-23.

⁹ Tsertsvadze A, Clar C, Court R, Clarke A, Mistry H, Sutcliffe P. Cost-effectiveness of manual therapy for the management of musculoskeletal conditions: a systematic review and narrative synthesis of evidence from randomized controlled trials. *J Manipulative Physiol Ther.* 2014 Jul-Aug;37(6):343-62.

¹⁰ Willich, S.N., et al., Cost-effectiveness of acupuncture treatment in patients with chronic neck pain. *Pain*, 2006. 125(1-2): p. 107-13.

¹¹ Haldeman S, Carroll L, Cassidy JD, Schubert J, Nygren A, Cn B, Joint Decade – Task Force on Neck P, Its Associated D: The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: executive summary. *Spine (Phila Pa 1976)* 2008, 33(4 Suppl):S5-7.

¹² Langenfeld A, Humphreys BK, Swanenburg J, Peterson CK. Prognostic factors for recurrences in neck pain patients up to 1 year after chiropractic care. *J Manipulative Physiol Ther.* 2015;38(7):458-64.

- ¹³ Council on Chiropractic Guidelines and Practice Parameters. Chiropractic management of neck related disorders. (accessed 9/16/2019 at http://www.ccgpp.org/neck_related_disorders.pdf)
- ¹⁴ Society for Acupuncture Research. Acupuncture for the treatment of cervical neck pain an evidence-based assessment. July 2017. (accessed 9/16/2019 at <http://www.asacu.org/wp-content/uploads/2017/09/Acupuncture-Evidence-Project-The.pdf>)
- ¹⁵ Gross A, et al. Manipulation or mobilisation for neck pain. *Cochrane Database Syst Rev*. 2010 Jan 20;(1):CD004249.
- ¹⁶ Leininger BD, Evans R, Bronfort G. Exploring patient satisfaction: a secondary analysis of a randomized clinical trial of spinal manipulation, home exercise, and medication for acute and subacute neck pain. *J Manipulative Physiol Ther*. 2014;37(8):593-601.
- ¹⁷ Gorrell LM, Beath K, Engel RM. Manual and Instrument Applied Cervical Manipulation for Mechanical Neck Pain: A Randomized Controlled Trial. *J Manipulative Physiol Ther*. 2016;39(5):319-29.
- ¹⁸ Gross A, Langevin P, Burnie SJ, et al. Manipulation and mobilisation for neck pain contrasted against an inactive control or another active treatment. *Cochrane Database of Systematic Reviews* 2015, Issue 9. Art. No.: CD004249.
- ¹⁹ Haavik H, Niazi IK, Holt K, Murphy B. Effects of 12 Weeks of Chiropractic Care on Central Integration of Dual Somatosensory Input in Chronic Pain Patients: A Preliminary Study. *J Manipulative Physiol Ther*. 2017;40(3):127-138.
- ²⁰ Coulter ID, Crawford C, Vernon H, et al. Manipulation and Mobilization for Treating Chronic Nonspecific Neck Pain: A Systematic Review and Meta-Analysis for an Appropriateness Panel. *Pain Physician*. 2019;22(2):E55-E70.
- ²¹ Moustafa IM, Diab AA, Taha S, Harrison DE. Addition of a Sagittal Cervical Posture Corrective Orthotic Device to a Multimodal Rehabilitation Program Improves Short- and Long-Term Outcomes in Patients With Discogenic Cervical Radiculopathy. *Arch Phys Med Rehabil*. 2016;97(12):2034-2044.
- ²² Yang JD, Tam KW, Huang TW, Huang SW, Liou TH, Chen HC. Intermittent Cervical Traction for Treating Neck Pain: A Meta-analysis of Randomized Controlled Trials. *Spine (Phila Pa 1976)*. 2017 Jul 1;42(13):959-965. doi: 10.1097/BRS.0000000000001948.
- ²³ Bryans R, Decina P, Descarreaux M, et al. Evidence-based guidelines for the chiropractic treatment of adults with neck pain. *J Manipulative Physiol Ther*. 2014 Jan;37(1):42-63.
- ²⁴ Maier M, Bronfort G, Evans R, et al. Manipulative therapy and exercise for seniors with chronic neck pain. *Spine J*. 2014 Sep 1;14(9):1879-89. Epub 2013 Nov 10.
- ²⁵ Thoomes EJ. Effectiveness of manual therapy for cervical radiculopathy, a review. *Chiropr Man Therap*. 2016;24:45.
- ²⁶ Bussi eres AE, Stewart G, Al-zoubi F, et al. The Treatment of Neck Pain-Associated Disorders and Whiplash-Associated Disorders: A Clinical Practice Guideline. *J Manipulative Physiol Ther*. 2016;39(8):523-564.e27.
- ²⁷ Araujo FX, Scholl schell M, Ribeiro DC. Effectiveness of Physiotherapy interventions plus Extrinsic Feedback for neck disorders: A systematic review with meta-analysis. *Musculoskelet Sci Pract*. 2017;29:132-143.
- ²⁸ De meulemeester KE, Castelein B, Coppieters I, et al. Comparing Trigger Point Dry Needling and Manual Pressure Technique for the Management of Myofascial Neck/Shoulder Pain: A Randomized Clinical Trial. *J Manipulative Physiol Ther*. 2017;40(1):11-20.
- ²⁹ Kojidi MM, Okhovatian F, Rahimi A, Baghban AA, Azimi H. Comparison Between the Effects of Passive and Active Soft Tissue Therapies on Latent Trigger Points of Upper Trapezius Muscle in Women: Single-Blind, Randomized Clinical Trial. *J Chiropr Med*. 2016;15(4):235-242.
- ³⁰ Sherman KJ, Cook AJ, Wellman RD, et al. Five-week outcomes from a dosing trial of therapeutic massage for chronic neck pain. *Ann Fam Med*. 2014;12(2):112-20.
- ³¹ Chow RT1 Johnson MI, Lopes-Martins RA, Bjordal JM. Efficacy of low-level laser therapy in the management of neck pain: a systematic review and meta-analysis of randomised placebo or active-treatment controlled trials. *Lancet*. 2009 Dec 5;374(9705):1897-908. doi: 10.1016/S0140-6736(09)61522-1. Epub 2009 Nov 13.
- ³² Becker JJ, Copeland SL, Botterbusch EL, Cohen RG. Preliminary evidence for feasibility, efficacy, and mechanisms of Alexander technique group classes for chronic neck pain. *Complement Ther Med*. 2018;39:80-86.
- ³³ Hurwitz EL, Carragee EJ, van der VeG, et al. Treatment of neck pain: noninvasive interventions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)*. 2008;33(4 Suppl):S123-52.
- ³⁴ Martin-Gomez C, Sestelo-Diaz R, Carrillo-Sanjuan V, Navarro-Santana MJ, Bard on-Romero J, Plaza-Manzano G. Motor control using cranio-cervical flexion exercises versus other treatments for non-specific chronic neck pain: A systematic review and meta-analysis. *Musculoskelet Sci Pract*. 2019 Jul;42:52-59. doi: 10.1016/j.msksp.2019.04.010. Epub 2019 Apr 20. PMID: 31030111.

-
- ³⁵ MacPherson, H., et al., Alexander Technique Lessons or Acupuncture Sessions for Persons With Chronic Neck Pain: A Randomized Trial. *Ann Intern Med*, 2015. 163(9): p. 653-62.
- ³⁶ Liang, Z., et al., Assessment of a traditional acupuncture therapy for chronic neck pain: a pilot randomised controlled study. *Complement Ther Med*, 2011. 19 Suppl 1: p. S26-32.
- ³⁷ Cameron, I.D., E. Wang, and D. Sindhusake, A randomized trial comparing acupuncture and simulated acupuncture for subacute and chronic whiplash. *Spine (Phila Pa 1976)*, 2011. 36(26): p. E1659-65.
- ³⁸ Chen, L., et al., Optimized acupuncture treatment (acupuncture and intradermal needling) for cervical spondylosis-related neck pain: a multicenter randomized controlled trial. *Pain*, 2021. 162(3): p. 728-739.
- ³⁹ Fu, W.B., et al., Analysis on the effect of acupuncture in treating cervical spondylosis with different syndrome types. *Chin J Integr Med*, 2009. 15(6): p. 426-30.
- ⁴⁰ White, A.R. and E. Ernst, A systematic review of randomized controlled trials of acupuncture for neck pain. *Rheumatology (Oxford)*, 1999. 38(2): p. 143-147.
- ⁴¹ Trinh, K., et al., Acupuncture for neck disorders. *Spine (Phila Pa 1976)*, 2007. 32(2): p. 236-243.
- ⁴² Fu, L.M., J.T. Li, and W.S. Wu, Randomized Controlled Trials of Acupuncture for Neck Pain: Systematic Review and Meta-Analysis. *J Altern Complement Med*, 2009. 15(2): p. 133-145.
- ⁴³ Jo, H.R., et al., Comparative effectiveness of different acupuncture therapies for neck pain. *Medicine (Baltimore)*, 2022. 101(33): p. e29656.
- ⁴⁴ Fu, L.M., J.T. Li, and W.S. Wu, Randomized Controlled Trials of Acupuncture for Neck Pain: Systematic Review and Meta-Analysis. *J Altern Complement Med*, 2009. 15(2): p. 133-145.
- ⁴⁵ Trinh, K., et al., Acupuncture for neck disorders. *Cochrane Database Syst Rev*, 2006. 3: p. CD004870.
- ⁴⁶ Liang, Z., et al., Assessment of a traditional acupuncture therapy for chronic neck pain: a pilot randomised controlled study. *Complement Ther Med*, 2011. 19 Suppl 1: p. S26-32.
- ⁴⁷ MacPherson, H., et al., Alexander Technique Lessons or Acupuncture Sessions for Persons With Chronic Neck Pain: A Randomized Trial. *Ann Intern Med*, 2015. 163(9): p. 653-62.
- ⁴⁸ Vickers, A.J., et al., Acupuncture for Chronic Pain: Update of an Individual Patient Data Meta-Analysis. *J Pain*, 2018. 19(5): p. 455-474.
- ⁴⁹ Willich, S.N., et al., Cost-effectiveness of acupuncture treatment in patients with chronic neck pain. *Pain*, 2006. 125(1-2): p. 107-13.
- ⁵⁰ Wirth B, Humphreys BK, Peterson C. Importance of psychological factors for the recovery from a first episode of acute non-specific neck pain - a longitudinal observational study. *Chiropr Man Therap*. 2016;24:9.
- ⁵¹ Cherkin DC, Herman PM. Cognitive and Mind-Body Therapies for Chronic Low Back Pain and Neck Pain: Effectiveness and Value. *JAMA Intern Med*. 2018;178(4):556-557.
- ⁵² Li Y, Li S, Jiang J, Yuan S. Effects of yoga on patients with chronic nonspecific neck pain: A PRISMA systematic review and meta-analysis. *Medicine (Baltimore)*. 2019;98(8):e14649.
- ⁵³ Allende S, Anandan A, Lauche R, Cramer H. Effect of yoga on chronic non-specific neck pain: An unconditional growth model. *Complement Ther Med*. 2018;40:237-242.
- ⁵⁴ Girard J, Girard A. The effects of qigong on neck pain: A systematic review. *Complement Ther Clin Pract*. 2019;34:23-29.
- ⁵⁵ Framework for Measuring Impact.
<https://www.hbs.edu/socialenterprise/Documents/MeasuringImpact.pdf> (accessed 5/29/2019)
- ⁵⁶ Willich SN, Reinhold T, Selim D, et al. Cost-effectiveness of acupuncture treatment in patients with chronic neck pain. *Pain*. 2006;125(1-2):107-13.
- ⁵⁷ MacPherson H, Tilbrook H, Richmond S, et al. Alexander Technique Lessons or Acupuncture Sessions for Persons With Chronic Neck Pain: A Randomized Trial. *Ann Intern Med*. 2015;163(9):653-62.
-